R2Play is a return-to-play assessment designed with stakeholders to better reflect the multi-domain demands of sport.

Learn more about the R2Play assessment here:








Holland Blcorview
Kids Rehabilitation Hospital





# R2Play Development: Fostering User Driven Technology that Supports Return-to-Play Decision-Making Following Pediatric Concussion

Josh Shore<sup>a,b</sup>, Danielle DuPlessis<sup>a,b</sup>, Emily Lam<sup>a</sup>, Fanny Hotze<sup>a</sup>, Ajmal Khan<sup>a</sup>, Setphanie McFarland<sup>a</sup>, Andrea Hickling<sup>a,c</sup>, Michael Hutchinson<sup>d</sup>, F. Virginia Wright<sup>a,b,e</sup>, Nick Reed<sup>a,b,d</sup>, Elaine Biddiss<sup>a,b,f</sup>, Shannon Scratch<sup>a,b,g</sup>

<sup>a</sup> Bloorview Research Institute, Holland Bloorview Kids Rehabilitation Hospital, Canada, <sup>b</sup> Rehabilitation Sciences Institute, University of Toronto, Canada, <sup>c</sup> Department of Occupational Science and Occupational Therapy, University of Toronto, Canada, <sup>d</sup> Faculty of Kinesiology and Physical Education, University of Toronto, Canada, <sup>e</sup> Department of Physical Therapy, University of Toronto, Canada, <sup>f</sup> Institute of Biomedical Engineering, University of Toronto, Canada, <sup>g</sup> Department of Pediatrics, University of Toronto, Canada.

## **Objective**

- Traditional return-to-play protocols after concussion rely on single-domain assessments (e.g., standing balance, running on a treadmill) and symptom self-reporting<sup>1</sup>
- These methods may fail to detect changes elicited by the *combined* physical, cognitive, and emotional demands of sport<sup>2,3</sup>
- To address this gap, we developed the *R2Play* system to facilitate a multi-domain return-to-play assessment


#### Methods

A design-thinking approach to development was used, involving:

- 1. Problem definition and early ideation via a scoping review and structured brainstorming
- 2. Needs-assessment interviews with stakeholders (6 clinicians and 4 youth sports coaches)
- 3. Building a prototype of R2Play and conducting usability testing via cognitive walkthroughs with 5 clinicians

## The R2Play Prototype

After problem definition and ideation, the prototype consisted of a tablet-button system that displays numbers and letters, and a clinician tablet that controls the assessment.



During the task, athletes run in a zig-zag pattern in an embodied *Trail Making Task* by pressing tablets in alphanumeric order.

# **Insights from Needs-Assessment Interviews**

- Interviews were analyzed using conventional content analysis
- A change table was constructed, in which the themes from user feedback were mapped to potential changes to the *R2Play* prototype

#### **Examples of Implemented Changes**

Category	Description	Change
Accessibility	Adapting R2Play for wheelchair users	Moved tablets onto elevated stands
Task	Navigating self in relation to moving/changing stimuli	Implemented scramble condition in which nodes change places during the trail
Interface	Ability to display results and use them to communicate with athletes	A graphical summary of results was developed with young athletes in mind

## **Usability Testing Results**

- The interface achieved a System Usability Scale score of 81% (SD=8.02), indicating "good" to "excellent" usability<sup>4</sup>
- Participants were comfortable navigating the interface and found the "flow" easy to follow

### Conclusion

- R2Play aligns with current best practice guidelines for return-to-play by simultaneously integrating physiological and neuropsychological measures across multiple domains
- With further testing and refinement, *R2Play* may provide clinicians with richer clinical data for making return-to-play decisions
- 1. McCrory, P. et al. Consensus statement on concussion in sport—the 5th international conference on concussion in sport held in Berlin, October 2016. Br J Sports Med -2017-097699 (2017) doi:10.1136/bjsports-2017-097699.
- 2. Fino, P. C. et al. Detecting gait abnormalities after concussion or mild traumatic brain injury: A systematic review of single-task, dual-task, and complex gait. *Gait & Posture* 62, 157–166 (2018).
- 3. Toong, T. *et al.* Sensitivity and Specificity of a Multimodal Approach for Concussion Assessment in Youth Athletes. *J Sport Rehabil* 1–10 (2021) doi:10.1123/jsr.2020-0279.
- 4. Sauro, J. & Lewis, J. R. *Quantifying the User Experience: Practical Statistics for User Research*. (Morgan Kaufmann, 2016).